Selected Abstracts From the Wilderness Medical Society's 2013 Winter Scientific Conference

Investigation into Recurrent Norovirus Outbreaks in Grand Canyon: Summer 2012

Background.—Norovirus is a highly contagious virus and the leading cause of acute gastrointestinal illness (AGI) in the United States. From 1994 to 2011, there were 8 norovirus outbreaks on Colorado River trips through Grand Canyon National Park. Only 3 of these outbreaks had identified sources (river water or food contamination).

Objective.—To determine the infection incidence, infectious agent, and possible contamination sources of norovirus on river trips during the 2012 summer season.

Methods.—Confidential illness reports were completed by AGI symptomatic persons and included date and location of illness onset and recovery. Toilet samples of collective fecal matter, a sample of vomitus, surface swabs of rafting equipment, and samples of toilet and water stops along the hiking corridor were collected and analyzed using RT-qPCR to detect the presence of norovirus.

Results.—From May 9 to September 28, 2012, 98 rafters from 10 different trips became ill with symptoms of AGI from norovirus. Attack rates on individual trips varied from 6% to 88%. Outbreaks occurred in 3 distinct clusters that involved 2 different strains of norovirus. All available toilet samples were positive for norovirus strains: 1 with the GI strain and 4 with the more common GII strain. The vomitus sample tested positive for GI. One of 12 fomite samples from rafting equipment was positive for GII norovirus. Swab samples collected along the hiking trail to the half-way point were negative for norovirus.

Conclusions.—Norovirus strains from the summer of 2012 outbreaks in Grand Canyon were not from a shared source such as contaminated food or river water. The results suggest that the virus was introduced by ill or asymptomatic persons actively shedding norovirus in their feces, and spread within, or between, river trips by different modes of transmission. This study reinforces the importance of education regarding prevention of norovirus and containment after outbreak.

Anne Magill-Collins PA
James D. Stoehr, PhD
Glendale, AZ, USA

Marlene Gaither, RS, MPA, ME
Flagstaff, AZ, USA

Charles P. Gerba, PhD
Tucson, AZ, USA

Acclimatization Practices of High-Altitude Recreationalists

Background.—Acute mountain sickness (AMS) will occur if ascent to high altitude is rapid. Any type of travel to altitudes around 2500 m (8000 feet) is associated with an increased risk of AMS. Prevention of AMS through acclimatization procedures such as gradual ascent and pharmacologic agents that aid in the body's accommodation to the hypobaric hypoxia resulting from high altitude exposure are key strategies. Forty percent of South Pole Station scientists and support staff during the 2006 and 2007 summer deployments chose to defer acclimatization medications before rapid ascent from McMurdo Station at sea level to the high altitude polar plateau at 2835 m (9300 feet), placing them at risk for having AMS (Anderson et al, 2011). Medical evacuation due to AMS is resource intensive and life-threatening to the person and to the rescue team. It also results in unsuccessful job performance for the person, precluding future polar plateau and frequently Antarctic deployment. The rationale and reasons for deferment of acclimatization guidelines is unknown.

Methods.—A convenience sample consisted of 120 volunteer high altitude recreationalists or backpackers, 18 years of age or older. A questionnaire was completed by subjects that included demographic information such as age, sex, altitude of residence, means of transport to calculate the rate of ascent, and the number of hours taken to acclimatize to the altitude at the trailhead. Acclimatization prophylaxis activities were documented, including the use of medications and personal home remedies. Baseline vital signs, height, weight, and body mass index were recorded for each subject. The instrument queried the volunteer about the existence of current AMS symptoms by completion of a Lake Louise Symptom Score questionnaire.

Results.—Despite recent Wilderness Medical Society (WMS) recommendations (Luks et al, 2010) stressing gradual ascent and medication prophylaxis for the prevention of AMS, the majority (55%) of subjects ($n = 66$) did not perceive the need for medications. Only 3 subjects (2.5%) were taking acetazolamide and only 4% had consulted a health professional before their high altitude sojourn. Sixty-one percent of the subjects took less than 24 hours to ascend to 9300 feet to begin their journey. Forty percent of the respondents camped at the trailhead or above 6800 feet the night before their departure, and 7 subjects (6%) took hikes to enhance acclimatization the day before their departure. Many subjects were veteran backpackers and reported their acclimatization symptoms could be controlled with hydration (12%), adequate rest, and anti-inflammatory drugs (40%) to control headaches,