Using a randomized double-blind design, 7 experienced male runners with no history of asthma inhaled either placebo, salbutamol 800 µg, or salbutamol 1600 µg 15 minutes before a 5-km time trial in either temperate (18°C) or hot (30°C) environments. Measured outcomes for each trial included overall time, heart rate, oxygen consumption, carbon dioxide production, minute ventilation, rate of perceived exertion, serum lactate, and urinary salbutamol concentration.

The investigators found no difference between placebo, salbutamol 800 µg, or salbutamol 1600 µg in either temperate or hot environments for overall time and all other outcome measures. Additionally, no adverse effects were detected with the use of salbutamol. Study limitations include sample size, variability of amount inhaled through pocket chamber, and body habitus variability. The authors conclude that current WADA guidelines are sufficient to avoid pharmaceutical performance enhancement.

(*J Sports Sci Med.* 2014;13:271–279). J Dickinson, J Hu, N Chester, et al.

Prepared by Matthew Stewart, University of Utah Global Medicine Fellow, Salt Lake City, UT.

SCIENTIFIC REPORTS

Diagnosis of Snake Envenomation Using a Simple Phospholipase A₂ Assay

Snake envenomations are a major public health concern worldwide, with an estimated 1 to 2 million cases reported annually and as many as 100,000 deaths per year. The accurate diagnosis of a snake envenomation requires the combination of patient history, clinical features, and laboratory testing. Unfortunately, resource-poor areas may lack trained medical personnel and the appropriate tests to accurately diagnose snake envenomation. The goal of this investigation was to evaluate a simple bedside test for a common component of snake venom and its ability to accurately predict envenomation.

Phospholipase A₂ (PLA₂) is a common toxin in snake venoms in nearly all elapid and viperid snakes. The authors recognize that although PLA₂ may not represent clinically important venom toxicity, it can be used to identify systemic envenomation. This prospective study included patients from 2 hospitals in Sri Lanka presenting with envenomation from a Russell's viper, hump-nosed pit viper, cobra, and krait snakes from 2007 to 2011. The PLA₂ activity and its correlation to a venom-specific assay was tested before and after antivenom administration and was compared to PLA₂ activity in nonenvenomated patients. The researchers found that PLA₂ activity directly correlated with free venom concentrations, and its activity decreased correspondingly with antivenom administration.

The authors propose that a bedside PLA₂ assay would be beneficial in areas with limited access to medical care. This assay could help to quickly identify patients with systemic

envenomation requiring antivenom administration. Limitations of the study include the small number of patients evaluated and envenomation of only 3 snake species tested with the assay. Additionally, the authors recognize the need to develop a PLA₂ assay that is less expensive and allows for individual patient testing.

(*Sci Rep.* 2014;4:4827). K Maduwage, MA O'Leary, G Isbister.

Prepared by Matthew Stewart, University of Utah Global Medicine Fellow, Salt Lake City, UT.

WORLD JOURNAL OF ORTHOPEDICS

Feet Injuries in Rock Climbers

Injuries to the upper extremities in rock climbers are widely discussed in current literature, but there is scarce literature concerning lower extremity injuries. Nearly 50% of acute injuries involve the legs and feet, most frequently strains, contusions, and fractures of the calcaneus and talus. The authors reviewed current literature and reported common lower extremity injury patterns and factors associated with injury. Wall-collision falls (climber impacts in a vertical plane) and ground falls (climber impacts in a horizontal plane) are the most common inciting factors for lower extremity injuries. Wall-collision falls are associated with contusions and compound fractures whereas ground falls can cause serious fractures of the talus, calcaneus, or ankle joint. Distortions and ligamentous injuries to the ankle are also seen in ground falls. Standard orthopedic guidelines suggest operative repair for displaced fractures as well as calcaneus or talus fractures in young, active patients.

A case of posttraumatic osteochondritis dissecans with a large osseous defect at the medial talar shoulder was discussed. This patient had failed a drill-hole arthroscopic procedure and subsequently had success with an osteograft with cartilage transplant. Three cases of peroneal tendon dislocation that underwent successful surgical management were also described.

Climbing foot injuries result from wearing climbing shoes that are unnaturally shaped or too small. Common problems are callosity, nail bed infections, pressure marks, neurologic complaints, claw toes, blisters, and subungual hematomas. A long-term problem is the development of a hallux valgus deformity. Climbers should be counseled not to wear extremely tight shoes, and future climbing shoes should adjust the biomechanical forces on the foot to prevent these problems. Limitations of this article include discrepancies in the existing literature on the incidence of lower extremity injuries in climbers.

(*World J Orthop.* 2013;4:218–228). V Schöffl, T Küpper.

Prepared by Pearly Ng, MD, University of Utah Wilderness Medicine Fellow, Salt Lake City, UT.