¹Department of Anesthesiology and Intensive Care Medicine, Medical University Hospital Innsbruck, Innsbruck, Austria, ²Austrian Board of Alpine Safety, Innsbruck, Austria, ³Alpine Police, Federal Ministry of the Interior, Vienna, Austria, ⁴Barts Heart Centre, St. Bartholomew's Hospital, Queen Mary University of London, London, UK

Introduction.—Since November 1, 2005, the Austrian Alpine Police records data for every mountain accident, which is reported to an emergency dispatch center, in a digital database. Still, no medical data on lightning injuries in the Austrian mountains are available.

Objective.—To analyze lightning casualties in the Austrian mountains with respect to accident mechanism, injury pattern, and medical outcome.

Methods.—The database was evaluated regarding emergencies caused by lightning that occurred from November 1, 2005, to December 31, 2014. Additionally, data for lightning strike victims who were treated at the Innsbruck University Hospital in Western Austria were included. This tertiary level hospital in the Eastern European Alps has a catchment area of approximately 25,000 km² and 3 million inhabitants. Data included sex, age, victim's activity at the time of the injury, severity of injury, and time and location of the accident.

Results.—In the study period, 67,807 alpine accidents were registered in the database, and 70,675 casualties were involved. The search for lightning casualties revealed 96 persons. Sixty-four of these fulfilled inclusion criteria for this study. Sixty of the victims (94%) survived the lightning accident. Four people died; among these were 2 hunters who were killed while in a high seat and a ranger who was felling trees. The fourth person was a hiker who was hiding from the thunderstorm below a mountain peak when he was hit.

Conclusions.—Most lightning accidents happen in late afternoon in summer. Climbers, mountaineers, and hikers are most commonly hit; however, they also make up the highest share of people who are active in the mountains. People next to trees seem to be at high risk of being struck by lethal lightning. Lightning accidents affecting humans should be better interlinked with hospital data to gain better insight into accident mechanism and outcome.

Trauma Training for Journalists: Developing and Implementing an Intercultural Medical Curriculum for Journalists Working in Regions of Conflict

David S. Young, MD, MS^{1,2}; Michael Puntis, MB, BChir³

¹University of Colorado Denver School of Medicine, Aurora, CO, USA,

³Rush University Medical Center, Chicago, IL, USA, ²St. George's University, London, UK

Introduction.—Almost 1200 journalists have died on assignment since 1992. Most deaths resulted from trauma (automobile accidents, gunfire) and occurred in the Middle East, Africa, and Latin America. While deaths of high-profile Western reporters have brought attention to inherent risks of reporting, 88% of deaths were local journalists earning a meager wage. In recognition of these dangers, major news agencies have endorsed freelance protection standards including basic medical training. However, trainings typically occur in North America and Western Europe—never reaching those most at risk.

Objective.—To establish and evaluate a culture-neutral course, to be delivered with equal effectiveness in a number of geographical areas, that will educate journalists on the basics of medical care in remote and resource-limited locations.

Methods.—A multinational team of physicians and paramedics was formed and led by a retired professional journalist. The team developed the curriculum based on wilderness medicine and combat experience. The course was administered in Beirut, chosen for its proximity to conflict and large media outlets. At the completion, students were tested on content and provided feedback.

Results.—Seventeen students of various nationalities (Lebanese, American, British, French, Canadian, Egyptian, Syrian, and Spanish), started and completed the course. Those speaking primarily English scored 80% on average compared with 71% for those speaking English secondarily and 60% for Arabic-speaking students.

Conclusions.—Developing and implementing a culture-neutral course proved to be a lengthy and challenging process. Live interpreters were invaluable for Arabic-speaking students, in addition to eliminating medical jargon and focusing on active demonstrations and scenarios. Despite this, language barriers

contributed to decreased retention of approximately one-quarter of the lecture material. Special attention to sex-norms in an Arabic country was necessary when undergoing scenario training. Lessons learned in the process will be applied to future TTJ courses and shared with those developing similar curricula.

Telesound: Can Real-Time Video Streaming of Ultrasound Imaging from Remote Locations Yield An Accurate Diagnosis?

Rachel E. Whitney; Antonio Riera; Lei Chen

Yale University, New Haven, CT, USA

Background.—Ultrasound has been proven to help medical decision-making in remote locations. Use of cell phones in medical care has become a popular trend given their availability, and 85% of the world has access to a 3G network.

Objective.—To evaluate live-stream video for interpretation of ultrasound using a 3G network.

Methods.—We evaluated the inter-rater reliability of assessment of ultrasound scans for abdominal free fluid, intussusception, and hip effusions. Deidentified 6-second video clips in .mp4 format were downloaded to a laptop: 58 focused assessment with sonography in trauma (FAST) scans, 44 scans evaluating for intussusception, and 28 scans evaluating for hip effusions. A bedside ultrasound expert evaluated each scan and marked it positive or negative for pathology. Using an iPhone 3G for remote “telesound” consultation, we used Skype to connect to our second ultrasound expert located in Ethiopia, who repeated the process over a live-stream video connection. When the remote expert returned from Ethiopia, evaluation of the scans was repeated as a bedside reviewer. Inter-rater reliability analysis using the Kappa statistic and percent agreement were performed to determine consistency among raters.

Results.—The Kappa values between the bedside and remote evaluators for FAST, intussusception, and hip effusion were 0.748 (95% CI 0.576–0.92), 0.816 (95% CI 0.648–0.984), and 0.764 (95% CI 0.519–1.0), respectively. The percent agreements after adjusting for chance were 86%, 80%, and 88%, respectively. The Kappa values between the bedside assessments by each evaluator were 0.851 (95% CI 0.71–0.992), 0.8 (95% CI 0.616–0.984), and 0.747 (95% CI 0.479–0.884), respectively. The percent agreements after adjusting for chance were 92%, 81%, and 88%, respectively.

Conclusion.—Live-stream video conference using 2 cellphones over a 3G network is a feasible and accurate method for consultation of ultrasound scans from a remote location.

CPR in Whitewater Drowning Victims: Is It Possible To Perform Adequate Chest Compressions Through a Personal Flotation Device (PFD)?

Scott McCorvey¹; Erin Lurie²; Robert D'Zio²; Michael Caudell²

¹Maine Medical Center, Portland, ME, USA, ²Medical College of Georgia at Augusta University, Augusta, GA, USA

Introduction.—Whitewater activities such as rafting, canoeing, and kayaking are growing in popularity throughout the country. More than 17 million Americans participated in these activities in 2008 alone. With this increase in participation has come an increase in paddling-related fatalities. It is estimated that 25% to 30% of all deaths are attributable to cardiac arrest.

Objective.—To examine the effects of a personal flotation device (PFD) on cardiopulmonary resuscitation (CPR) chest compression depth.

Methods.—This is a prospective study, using volunteers certified in CPR by the American Heart Association (AHA) and the Laerdal Medical Simulation Mannequin (SimMan) to evaluate the efficacy of chest compressions while wearing 3 different PFD styles. Compression depth, rate, and variance were recorded during 2-minute cycles of chest compressions and averaged for each test group. Statistical analysis was performed using a student's t-test.

Results.—Data were compared between the control group and 3 different PFD styles measuring compression depth and rate (n = 37 per group). All groups had adequate rates of compression. The mannequin without a PFD (control) demonstrated an average depth of 49.5 mm. The 3 PFD styles allowed an average compression depth of 48.7 mm (P = .61), 45.7 mm (P < .016), and 45.2 mm (P < .019).

Conclusion.—2015 AHA guidelines recommend adult CPR compression depth of 50 to 60 mm. Other studies have shown maximum survival at lower